


# shortRNA: A flexible framework for the analysis of short RNA sequencing data applicable to studies on epigenetic inheritance

Pierre-Luc Germain<sup>†1,2,3,4</sup>, Deepak K. Tanwar<sup>†1,2,3</sup>, Isabelle M. Mansuy<sup>1,2</sup>



<sup>1</sup> Institute for Neuroscience, Department of Health Sciences and Technology, ETH Zürich  
<sup>2</sup> Laboratory of Neuroepigenetics, Brain Research Institute, Medical Faculty of the University of Zürich and Institute for Neuroscience, Department of Health Sciences and Technology of the ETH Zürich  
<sup>3</sup> Statistical Bioinformatics Group, Swiss Institute of Bioinformatics  
<sup>4</sup> Statistical Bioinformatics Group, University of Zürich

## Introduction

Short RNAs, categorized as non-coding RNA molecules, are less than 200 nucleotides in length and play a vital role in the regulation of the genome. They have been implicated in a large number of biological processes and pathological conditions. Moreover, they are shown to be altered in different models of epigenetic inheritance [1–3]. Short RNA studies are hampered by a number of technical issues, including the bioinformatic analysis of short RNA sequencing data:

1. Analysis methods are not simultaneously optimized towards all known short RNA types (e.g. miRNA, piRNA, rasiRNA, siRNA, snoRNA, tsRNA, tRFs, srRNA and U-RNA). On top of multiplying the work needed for an extensive analysis of the data, this can potentially create misassignment mistakes.
2. Current methods either do not deal adequately with post-transcriptional modifications (for genome-based methods); or if they do (transcript-based methods), they do not deal with unannotated features.
3. Current methods do not adequately account for the hierarchical organization of the features one might want to quantify or test.
4. There is still no consensus on the most appropriate normalization method for short-RNA-Seq data.

### Example 1

#### The issue of post-transcriptional modifications:

tRNA-iMet-CAT-4 is transcribed from chrX, and as other tRNAs receives post-transcriptionally the 3' addition of the nucleotides CCA:

... TCCTCTGCTT → aligns to multiple locations on the genome  
 ... TCCTCTGCTTCCA → does not align to the genome

To deal with this, researchers have often simply trimmed 3' CCAs before alignment, however this can often results in the read becoming ambiguous when instead it initially wasn't, as in the case above. One solution to this issue is to build a custom genome that is complemented with known, post-transcriptionally modified transcripts [4].

### Example 2

#### The issue of ambiguity between related features:

tRNA typically have several copies across the genome - for instance tRNA-Ala-AGC has 23 nearly-identical copies. While a genome alignment will make most reads from such features ambiguous (i.e. multimapping), from a functional point of view it is irrelevant from which exact location they came from. This issue becomes even more critical with tRNA fragments, which often have conserved sequences across different tRNAs. One way to address this issue is to aggregate reads into functional equivalence classes, i.e. higher level than specific genes/transcripts.



## Objectives

Because of these shortcomings, we developed a new analysis framework that addresses these issues using alternative nested equivalence classes over a customized annotation. We present this approach and package, and show how it can be used to redress biases in the quantification of both specific RNA as well as large RNA classes.

## Methods

shortRNA R package is an extension to the TreeSummarised-Experiment object.

### Database sources




Figure 1: Databases used for mouse.

## Tree structure




Figure 2: An example tree structure from tRNAs.

## Pipeline for short RNA-Seq data analysis




Figure 3: Pipeline for data analysis.

## Results

Tree-based read assignment recover a large fraction of reads considered ambiguous from a mapping point of view




Figure 4: Reads assignment (Sperm short RNA-Seq [3]).

## Transcripts abundance plot for uniquely mapping fragments




Figure 5: Unique fragments abundance for a transcript (example).

## RNA biotype and their proportions




Figure 6: Read proportions from samples (example).

## Comparison of shortRNA with other methods

We compared 20 published tools for 21 features.

Features	ncPro-Seq	SPORTS 1.0	exceRpt	RAPID	shortRNA
Languages	R, Perl, Python	Perl, R	Java, R	R, Bash, Perl	R
Quality control	✓	✓	✓	✓	✓
Alignment/ Mapping	Bowtie	✓	✓	✓	✓
Databases	miRBase, Rfam, RepeatMasker, User defined	miRBase, rRNA (NCBI), GtRNadb, piRNA, Ensembl, Rfam	Gencode, miRBase, GtRNadb, circBase	User defined	Gencode, GtRNadb, miRBase, piRNA precursors, User defined
Normalization	✗	✗	✓	✓	✓
Differential analysis	✗	✗	✓	✓	✓
Functional analysis	✗	✗	NA	✗	✓
Exploratory data analysis	✓	✓	✓	✓	✓
Adequate handling of post-transcriptional modifications	✗	✓	✗	✗	✓
Unannotated transcripts/ Novel predictions	✓	✓	✓	✗	✓
Heirarchical	✗	✗	✗	✗	✓
isoMirs	✗	✗	✗	✗	✓
User interface	CLI, GUI	CLI	CLI, GUI	CLI	CLI, GUI
Implementation	Tool, Webserver	Tool	Tool, Webserver, Docker	Tool, Conda	R package

## Conclusion

- Standalone R package for short RNA-Seq data analysis.
- Extension to the TreeSummarisedExperiment object.
- QC, alignment, differential analysis, and functional analysis within package.
- Heirarchical: could easily be extended to additional feature trees (for example Vault RNAs).

## Code

Github: [mansuylab/shortRNA](https://github.com/mansuylab/shortRNA)

## References

1. Posner R, Toker IA, Antonova O, Star E, Anava S, Azmon E, et al. Neuronal small RNAs control behavior transgenerationally. Cell. 2019;177:1814–1826.e15. doi:10.1016/j.cell.2019.04.029.
2. Bohacek J, Mansuy IM. Molecular insights into transgenerational non-genetic inheritance of acquired behaviours. Nature Reviews Genetics. 2015;16:641–52. doi:10.1038/nrg3964.
3. Gapp K, Jawaid A, Sarkies P, Bohacek J, Pelczar P, Prados J, et al. Implication of sperm RNAs in transgenerational inheritance of the effects of early trauma in mice. Nature Neuroscience. 2014;17:667–9. doi:10.1038/nn.3695.
4. Hoffmann A, Fallmann J, Vilardo E, Mörl M, Stadler PF, Amman F. Accurate mapping of tRNA reads. Bioinformatics. 2017;34:1116–24. doi:10.1093/bioinformatics/btx756.